




# Clopixol

You are taking Clopixol due to feeling unwell
Clopixol comes as an injection, it will be injected into your bottom
When you take Clopixol you may: Feel stiff Gain weight Become confused Have blurred vision
See a doctor if you: Have difficulties breathing Itchy skin Swelling of the lips, face or tongue
Clopixol will help you feel better





#### **Haloperidol**








### Movicol

You are taking Movicol because you suffer from constipation.
Movicol comes in a powder that is mixed with water.
Movicol may make you: Feel sick Have tummy pains Have a sore bottom
See a doctor if: Have an itchy skin Have problems breathing
Movicol will help you to be able to go to toilet.





#### Carbamazepine







# Omeprazole

You are taking Omeprazole because you have indigestion or heart burn
Omeprazole comes as a tablet. Tablets should be swallowed with a glass of water
While taking Omeprazole you may: Feel sick Have a headache Feel tired Have tummy pains
See a doctor if: You have swelling of the lips, face of throat You have a fever Your skin blisters
Omeprazole will help to feel more comfortable






# Sodium valporate

You are taking Sodium Valporate because you have epilepsy
Sodium Valporate comes as a liquid, tablets or as granules. The tablets should be taken with a glass of water the granules should be sprinkled onto a small amount of soft food or into a cold drink.
While taking Sodium Valporate you may: Feel sick Have tummy pains Put on weight Feel faint
See a doctor if: You have a rash You face, lips, tongue or throat swell up You have problems breathing
Sodium Valporate will help to reduce the number of epileptic fits you have















You are taking Clomipramine because you have depression, or a phobia
Clomipramine comes as tablets. Tablets should be taken with water
While taking Clomipramine you may: Feel hungry Put weight on Feel sick Have problems to wee
See a doctor if: You have a face You have difficulty breathing
Clomipramine will help you to feel better






# Docusate Sodium

	You are taking Docusate Sodium because you are constipated or to stop you becoming constipated
	Docusate Sodium comes as a tablet. Tablets should be swallowed whole with a glass of water
	While taking Docusate Sodium you may: Feel sick Have tummy pains Have a rash
	See a doctor if any of the side effect last for more then a few days or become a problem
	Docusate Sodium will help you to go to the toilet















You are taking Melatonin because you have problems getting to sleep or staying asleep
Melatonin comes as tablets and should be taken with food 1-2 hours before you go to bed.  Tablets should be taken whole with a glass of water
While taking Melatonin you may: Become constipated Have tummy pains Feel sick
See a doctor if the symptoms get worse
Melatonin will help you to sleep



### Senna



You are taking Senna because you are constipated
Senna comes as tablets or as syrup. Tablets should be taken with water
While taking Senna you may: Lose weight Have diarrhoea Have constipation
See a doctor if any of the symptoms get worse
Senna will help you to be able to go to the toilet







You are taking Lactulose because you are constipated
Lactulose is a liquid
While taking Lactulose you may: Have diarrhoea Feel bloated
See a doctor if the side effects get worse
Lactulose will help you to go to toilet