
















## Olanzapine

	<p>You are taking Olanzapine because you are hearing, seeing or sensing things that are not really there and stopping activities you normally do</p>
	<p>Olanzapine comes as tablets. Tablets should be taken with water</p>
	<p>When you take Olanzapine you may:</p> <ul style="list-style-type: none"><li>Feel sleepy</li><li>Feel hot</li><li>Feel hungry</li><li>Feel weak</li><li>Feel stiff</li></ul>
	<p>See a doctor if you have a rash</p>
	<p>Olanzapine will help you feel calm and relaxed</p>






## Pregablin

	<p>You are taking Pregablin because of feeling anxious or because you have epilepsy</p>
	<p>Pregablin come as tablets. Tablets should be taken with water</p>
	<p>When you take Pregablin you may:</p> <ul style="list-style-type: none"><li>Feel hungry</li><li>Feel dizzy</li><li>Put on weight</li><li>Have a dry mouth</li></ul>
	<p>See a doctor if :</p> <ul style="list-style-type: none"><li>Your face or tongue swells up</li><li>Your skin is red, blisters or peels</li></ul>
	<p>Pregablin will help you feel calm and relaxed</p>






## Aripiprazol (Abilify)

	<p>You are taking Aripiprazol because you are hearing, seeing or sensing things that are not really there</p>
	<p>Aripiprazol comes as tablets. Tablets should be taken with water</p>
	<p>When you take Aripiprazol you may:</p> <ul style="list-style-type: none"><li>Feel dizzy or weak</li><li>Feel restless</li><li>Feel sick</li><li>Have problems sleeping</li><li>Have a headache</li><li>Become constipated</li></ul>
	<p>See a doctor if you:</p> <ul style="list-style-type: none"><li>feel stiff</li><li>feel shaky</li><li>have blurred vision</li><li>have a racing heart</li></ul>
	<p>Aripiprazol will help you feel calm and relaxed</p>






## Celebrex

	<p>You are taking Celebrex because you are in pain</p>
	<p>Celebrex comes as tablets. Tablets should be taken with water.</p>
	<p>When you take Celebrex you may: have a sore throat have tummy pains feel sick have problems sleeping</p>
	<p>See a doctor if: you have problems breathing your face swells up you have a rash</p>
	<p>Celebrex will help to ease the pain</p>






## Lorazepam

	You are taking Lorazepam because you are feeling anxious
	Lorazepam comes as tablets. Tablets should be taken with water.
	When you take Lorazepam you may: Feel unsteady Feel light-headed and faint Feel sleepy and slow Feel confused Have a headache
	See a doctor if you: Have a rash or blotches on your skin
	Lorazepam will help you to feel calm and relaxed






## Procyclidine

	You are taking Procyclidine because your other medications are making you feel stiff or shaky
	Procyclidine comes as tablets or liquid. Tablets should be taken with water
	When you take Procyclidine you may: Have blurred vision Have a dry mouth Have problems to wee Feel sick
	See a doctor if you: Have unusual body movements Have uncontrolled movements of the face and tongue
	Procyclidine will stop you feeling stiff or shaky

## Lithium






	<p>You are taking Lithium because you are feeling unwell</p>
	<p>Lithium comes as tablets or liquid. Tablets should be taken with water</p>
	<p>When you take Lithium you may:</p> <ul style="list-style-type: none"> <li>Need to drink more</li> <li>Need to wee more</li> <li>Feel confused</li> <li>Feel sick</li> <li>Feel sleepy</li> <li>Feel unsteady on your feet</li> </ul>
	<p>See a doctor if you:</p> <ul style="list-style-type: none"> <li>Have dizziness or blurred vision</li> <li>Hear ringing in your ears</li> <li>Feel weak or numb in your arms and legs</li> <li>Have sickness and diarrhoea</li> </ul>
	<p>Lithium will help you to feel better</p>

## Fluoxetine






	<p>You are taking Fluoxetine because you have depression or obsessive-compulsive disorder</p>
	<p>Fluoxetine comes as tablets or liquid. Tablets should be taken with water</p>
	<p>When you take Fluoxetine you may:</p> <ul style="list-style-type: none"><li>Have problems sleeping</li><li>Have an upset tummy</li><li>Feel sick</li><li>Feel confused</li><li>Need to wee more</li></ul>
	<p>See a doctor if you have:</p> <ul style="list-style-type: none"><li>A rash all over your skin</li><li>Itchy skin</li><li>A fever</li></ul>
	<p>Fluoxetine will help you feel calm and relaxed</p>



## Diazepam

	<p>You are taking Diazepam because of feeling anxious or you have epilepsy</p>
	<p>Diazepam comes as tablets and liquid. Tablets should be taken with water</p>
	<p>When you take Diazepam you may:</p> <ul style="list-style-type: none"><li>Feel unsteady</li><li>Feel light-headed and faint</li><li>Feel sleepy and slow</li><li>Feel confused</li><li>Have a headache</li></ul>
	<p>See a doctor if you:</p> <ul style="list-style-type: none"><li>Have a rash or blotches on your skin</li></ul>
	<p>Diazepam will help you to feel calm and relaxed</p>

## Risperidone

	<p>You are taking Risperidone Because you have a condition that affects the way you think, feel and act</p>
	<p>Risperidone comes as tablets or liquid. Tablets should be taken with water</p>
	<p>When you take Risperidone you may: Have a dry mouth Feel drowsy Feel restless Have problems sleeping</p>
	<p>See a doctor if you: Feel faint Have a rash Feel stiff or shaky Have a pounding heart</p>
	<p>Risperidone will help you feel calm and relaxed</p>